

Vacuum Polarization Effects in the Lorentz and PCT Violating Electrodynamics

G. Bonneau^{[a]*}, L. C. Costa^{[a][b]} and J. L. Tomazelli^{[a][c]}

^[a] *Laboratoire de Physique Théorique et Hautes Energies[†], Universités de Paris VI et Paris VII,
2 Place Jussieu, 75251 Paris Cedex 05, France.*

^[b] *On leave from Departamento de Física, Universidade Federal de São Carlos,
P.O. Box 676, 13565-905, São Carlos, SP, Brazil.*

^[c] *On leave from Departamento de Física e Química, Universidade Estadual Paulista,
Av. Dr. Ariberto Pereira da Cunha 333, 12500-000 Guaratinguetá, SP, Brazil.*

Abstract

In this work we report new results concerning the question of dynamical mass generation in the Lorentz and PCT violating quantum electrodynamics. A one loop calculation for the vacuum polarization tensor is presented. The electron propagator, “dressed” by a Lorentz breaking extra term in the fermion Lagrangian density, is approximated by its first order: this scheme is shown to break gauge invariance. Then we rather consider a full calculation, to second order in the Lorentz breaking parameter: we recover gauge invariance and use the Schwinger-Dyson equation to discuss the full photon propagator. This allows a discussion on a possible photon mass shift as well as measurable, observable physical consequences, such as the Lamb-shift.

PACS numbers : 11.10.Wx, 12.20.-m

**E-mail addresses* : bonneau@lpthe.jussieu.fr (G. Bonneau); lccosta@ift.unesp.br (L.C. Costa)

[†]Unité associée au CNRS UMR 7589.

I. INTRODUCTION

The PCT theorem states that if a field theory satisfies the following axioms: *a) locality, b) Lorentz invariance and c) analiticity of the Lorentz group representation in the boost parameters*, the PCT transformation is a symmetry of the theory itself [1]. In this context, the invariance under the Lorentz group represents one of the fundamental axioms in the construction of a relativistic quantum field theory, which includes the minimal $SU(3) \times SU(2) \times U(1)$ Standard Model.

The possibility of nature to reveal a small violation of the Lorentz and PCT symmetries has been object of an intense research which includes different areas of physics, ranging from Quantum Optics to Neutrino Physics [2]. However, up to now, there is no conclusive experimental evidence for the violation of axiom *b)*.

From the theoretical standpoint, there arose a controversy on a possible Chern-Simons-like term generated through radiative corrections in an extended version of Quantum Electrodynamics (QED) [3] - [8]. In fact, some authors claimed that the addition to the QED Lagrangian density of a renormalizable and Lorentz violating term such as *

$$\mathcal{L}_{LB} = -b^\alpha \bar{\psi} \gamma_\alpha \gamma_5 \psi \quad (1)$$

induces a Chern-Simons-like term, namely,

$$\mathcal{L}_{CS} = \frac{1}{2} c_\mu \epsilon^{\mu\alpha\beta\gamma} F_{\alpha\beta} A_\gamma. \quad (2)$$

*where :

- b^α is a set of four constants, which selects a preferential direction in space-time, thus violating Lorentz invariance;
- ψ represents the electron field;
- γ_5 a hermitian matrix with the properties $\{\gamma_5, \gamma_\alpha\} = 0$ and $\text{tr} \gamma_5 \gamma^\alpha \gamma^\beta \gamma^\gamma \gamma^\delta = 4i \epsilon^{\alpha\beta\gamma\delta}$.

thus violating the Lorentz and PCT symmetry. *Note that such a term (2) does not destroy the gauge invariance of the action.*

As a matter of facts, at least two questions arose :

- As soon as Lorentz symmetry is violated, one could wonder about the validity of any calculation : should one consider - as done for example in [5] - that the loop momenta phase space is no longer Lorentz covariant? Should the regularization preserve Lorentz invariance? Here, we shall follow the conservative point of view initiated by Colladay and Kostelecky in which the sole breaking of Lorentz invariance comes from the added Lagrangian (1).
- The Lagrangian (1) modifies the fermion propagator and two approaches are possible. Do an expansion in b^α of the propagator which amounts to consider (1) as a new interaction: we call this approach a perturbative one. Either try to compute with the complete b^α dependent fermion propagator: we call this approach a non-perturbative one.

Then, if the Lorentz breaking is the minimal one and the electron propagator expanded in powers of b^α , and if the theory is correctly defined through Ward identities and normalisation conditions, no Chern-Simons term appears [7][†]. A similar result was first obtained in the analysis due to Coleman and Glashow [3] and this result was afterwards confirmed through a proper-time approach by Sytenko and Rulik [8].

In this work we intend to reexamine the role of gauge invariance and the issue of its possible breaking; moreover, we want to study a possible dynamical mass generation for the photon. So, we compute the vacuum polarization amplitude $\Pi^{\mu\nu}$ in QED (regulated

[†]This is due to the fact that such a term (2) is a kind of minor modification of the gauge fixing term, and then no renormalised. Then, if absent at the classical level, it remains absent at the loop level.

by means of the gauge invariant Pauli-Villars-Rayski (P-V-R) scheme [10]) when the tiny Lorentz breaking term (1) is added.

First, in Section 2, and in the spirit of [13], we consider the loop expansion of the vacuum polarization tensor when the fermion propagator that results from the QED Lagrangian density modified by term (1) is the upper mentioned "non perturbative one". However, to simplify the discussion and thanks to the smallness of parameter b^α , we approximated the infinite series in b^α by its linear contribution, to each order in the $\hbar$ perturbative series. This procedure essentially consists in dressing the fermion propagator with linear corrections in b^α . However, this scheme suffers from many difficulties and we do not pursue its study.

Second, in a more standard way, in Section 3 we consider (1) as an interaction term and then do a double expansion in b_α and $\hbar$. We give the complete one loop calculation to second order in the breaking parameter b_α . As expected from general results [7] we check that gauge invariance is recovered and, if an adequate normalization condition is chosen, the photon may remain massless.

Then, in Section 4, the full photon propagator will be obtained by summing the perturbative series through the Schwinger-Dyson Equation [11], allowing for a detailed discussion on the dynamical mass generation (the pole shifting in the photon propagator). Consequences on the Lamb-shift are also addressed.

Finally, some concluding remarks are offered in Section 5.

II. THE ONE LOOP VACUUM POLARIZATION TENSOR IN EXTENDED QED

In the extended version of QED, defined by the Lagrangian density

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2\alpha}(\partial_\mu A^\mu)^2 + \bar{\psi}(i\not{\partial} - e\not{A} - \not{b}\gamma_5 - m)\psi, \quad (3)$$

the fermion propagator is

$$S(l) = \frac{i}{\not{l} - m - \not{b}\gamma_5} = \sum_{n=0}^{\infty} \frac{i}{\not{l} - m} \left\{ -i \not{b}\gamma_5 \frac{i}{\not{l} - m} \right\}^n = \sum_{n=0}^{\infty} S_n(l),$$

and its linear approximation, used in that section, writes:

$$S^L(l) = S_0(l) + S_1(l) = \frac{i}{\not{l} - m} - i \frac{i}{\not{l} - m} \not{l} \gamma_5 \frac{i}{\not{l} - m}, \quad (4)$$

In that linear approximation, the one-loop vacuum polarization tensor is

$$\Pi^{\mu\nu}(p, m, b) = -(-i e)^2 \int \frac{d^4 l}{(2\pi)^4} \text{Tr}[\gamma^\mu S^L(l) \gamma^\nu S^L(l + p)]. \quad (5)$$

$\Pi^{\mu\nu}(p, m, b)$ then admits the decomposition

$$\Pi^{\mu\nu}(p, m, b) = \Pi_0^{\mu\nu}(p, m) + \Pi_b^{\mu\nu}(p, m, b) + \Pi_{bb}^{\mu\nu}(p, m, b). \quad (6)$$

In the last expression,

$$\Pi_0^{\mu\nu}(p, m) = e^2 \int \frac{d^4 l}{(2\pi)^4} \text{Tr}[\gamma^\mu S_0(l) \gamma^\nu S_0(l + p)]$$

is the usual QED vacuum polarization tensor,

$$\Pi_b^{\mu\nu}(p, m, b) = e^2 \int \frac{d^4 l}{(2\pi)^4} \text{Tr}[\gamma^\mu S_0(l) \gamma^\nu S_1(l + p) + \gamma^\mu S_1(l) \gamma^\nu S_0(l + p)] \quad (7)$$

is the linear b_α contribution, and

$$\Pi_{bb}^{\mu\nu}(p, m, b) = e^2 \int \frac{d^4 l}{(2\pi)^4} \text{Tr}[\gamma^\mu S_1(l) \gamma^\nu S_1(l + p)] \quad (8)$$

is a $\mathcal{O}(b^2)$ contribution to the polarization tensor, whose thorough calculation will be one of the focuses of the present work.

Diagrammatically, the expansion (6) may be represented by the series

$$\begin{aligned} \Pi^{\mu\nu}(p, m, b) \equiv & \text{Diagram with two external wavy lines labeled } p, \text{ and a shaded oval with indices } \nu, \mu \text{ on the vertices} \\ & = \text{Diagram with two external wavy lines labeled } l+p, \text{ and a loop with vertices labeled } l \text{ and } l+p \\ & + \text{Diagram with two external wavy lines labeled } l+p, \text{ and a loop with vertices labeled } l \text{ and } l+p \text{ (crosses on vertices)} \\ & + \text{Diagram with two external wavy lines labeled } l+p, \text{ and a loop with vertices labeled } l \text{ and } l+p \text{ (crosses on vertices)} \\ & + \text{Diagram with two external wavy lines labeled } l+p, \text{ and a loop with vertices labeled } l \text{ and } l+p \text{ (crosses on vertices)} \end{aligned}$$

where

$$-\times- \equiv \frac{\text{Diagram with a vertical dashed line labeled } b_\alpha \text{ and a vertex labeled } i\gamma^\alpha \gamma_5}{i\gamma^\alpha \gamma_5} + \mathcal{O}(b^2)$$

represents the linear b_α insertion to the (internal) fermion lines.

By power counting, $\Pi_{bb}^{\mu\nu}(p, m, b)$, $\Pi_b^{\mu\nu}(p, m, b)$ and $\Pi_0^{\mu\nu}(p, m)$ are respectively logarithmically, linearly and quadratically divergent in the ultra-violet region. Of course, a regularization procedure is needed: in order to preserve as much as possible the symmetries of the classical theory (3) (*i.e.* gauge invariance), the Pauli-Villars-Rayski (P-V-R) regularization prescription will be employed [10]. In this scheme, auxiliary fermion masses satisfying specific conditions are introduced. The original theory is recovered at the end of the calculations, by taking arbitrary large values for the auxiliary masses.

Let us recall also that thanks to gauge invariance, the degree of divergence of the tensor $\Pi_0^{\mu\nu}(p)$, identical to the pure QED sector, is reduced from 2 to 0 (logarithmic divergence).

To be consistent with the P-V-R scheme, the tensor $\Pi^{\mu\nu}(p, m, b)$ must be regulated as a whole object, which implies the replacement of expression (6) by the sum

$$\Pi^{\mu\nu}(p, b) = \sum_{i=0}^N c_i (\Pi_0^{\mu\nu}(p, m_i) + \Pi_b^{\mu\nu}(p, m_i, b) + \Pi_{bb}^{\mu\nu}(p, m_i, b)), \quad (9)$$

where each term in the r.h.s. of (9) retains the original functional form, except the fact that $m \rightarrow m_i$ (note also that we consider the same Lorentz breaking parameter b^α for all fermions). By analysing the structure of (9), it can be shown that, for auxiliary masses satisfying the conditions:

$$\begin{aligned} a) \quad \sum_{i=0}^N c_i &= 0, \\ b) \quad \sum_{i=0}^N c_i m_i^2 &= 0, \end{aligned} \quad (10)$$

the divergences in $\Pi^{\mu\nu}(p, m, b)$ all disappear. In the above expressions, $c_0 = 1$ and $m_0 = m$ is the electron mass.

The results may be written as[‡]

$$\Pi_0^{\mu\nu}(p, m_i) = [g^{\mu\nu} p^2 - p^\mu p^\nu] \Pi_0(p^2, m_i), \quad (11)$$

[‡]Details on that calculation will be published elsewhere.

$$\Pi_b^{\mu\nu}(p, m_i, b) = \epsilon^{\mu\nu\alpha\beta} p_\alpha b_\beta \Pi_b(p^2, m_i), \quad (12)$$

$$\begin{aligned} \Pi_{bb}^{\mu\nu}(p, m_i, b) = & \mathbb{A}((p.b)^2, b^2, p^2, m_i) g^{\mu\nu} + \mathbb{B}(p^2, m_i) b^\mu b^\nu + \\ & + \mathbb{C}(p^2, m_i)(p.b)(b^\mu p^\nu + b^\nu p^\mu) + \mathbb{D}((p.b)^2, b^2, p^2, m_i) p^\mu p^\nu, \end{aligned} \quad (13)$$

where

$$\begin{aligned} \Pi_0(p^2, m_i) &= i \frac{e^2}{12\pi^2} \left\{ \log \frac{m_i^2}{m^2} - p^2 \int_0^1 dz \frac{[1 - 2Z - 8Z^2]}{2\Delta_i} \right\}, \\ \Pi_b(p^2, m_i) &= i \frac{e^2}{2\pi^2} \left\{ p^2 \int_0^1 dz \frac{Z}{\Delta_i} \right\}, \\ \mathbb{A}((p.b)^2, b^2, p^2, m_i) &= -i \frac{e^2}{\pi^2} \left\{ \frac{b^2}{6} \log \frac{m_i^2}{m^2} + \int_0^1 dz \left[\frac{-1 - 4Z + 8Z^2}{12\Delta_i} b^2 p^2 + \frac{Z^2}{\Delta_i} (b.p)^2 + \right. \right. \\ &\quad \left. \left. + \frac{Z^2}{2[\Delta_i]^2} p^2 [(b.p)^2 - b^2 p^2] \right] \right\}, \\ \mathbb{B}(p^2, m_i) &= -i \frac{e^2}{\pi^2} \left\{ -\frac{1}{6} \log \frac{m_i^2}{m^2} + p^2 \int_0^1 dz \left[\frac{1 + 10Z - 8Z^2}{12\Delta_i} + \frac{Z^2}{[\Delta_i]^2} p^2 \right] \right\}, \\ \mathbb{C}(p^2, m_i) &= -i \frac{e^2}{\pi^2} \left\{ \int_0^1 dz \left[-\frac{Z(1 - 2Z)}{\Delta_i} - \frac{Z^2}{[\Delta_i]^2} p^2 \right] \right\}, \\ \mathbb{D}((p.b)^2, b^2, p^2, m_i) &= -i \frac{e^2}{\pi^2} \left\{ \int_0^1 dz \left[\frac{3Z^2}{\Delta_i} b^2 + 2 \frac{Z^3}{[\Delta_i]^2} ((p.b)^2 + b^2 p^2) \right] \right\}. \end{aligned} \quad (14)$$

In those expressions, we have set

$$Z = z(1 - z),$$

$$\Delta_i = m_i^2 - z(1 - z)p^2.$$

Moreover, to simplify our results we used some integration by parts on z and the symmetry of the integral when z is changed into $(1 - z)$: this allows us to rewrite the integrands as functions of Z .

A few comments are in order :

- The scale in the logarithms has been (arbitrarily) chosen to be the electron mass, thanks to the condition (10-a) $\sum_{i=0}^{i=N} c_i = 0$.
- Since $\Pi_b^{\mu\nu}$ vanishes at $p^2 = 0$, no Chern-Simons-like term appear [7].

- The vacuum polarization tensor is no longer transverse as :

$$p_\nu \Pi^{\mu\nu} = [\mathbb{A} + (p.b)^2 \mathbb{C} + p^2 \mathbb{D}] p^\mu + (p.b)[\mathbb{B} + p^2 \mathbb{C}] b^\mu$$

and neither of the two square brackets vanish (just look at the logarithms). Moreover, no local counterterm can be added in order to recover transversality. Then gauge invariance is definitely broken in that scheme. This might be a consequence of our specific expansion in b_α and, indeed, in the scheme of section 3, we shall recover gauge invariance at the one loop level by introducing new missing Feynman graphs, which amounts to consider all possible permutations among photon and b_α linear insertions as, for example, in the case of the gauge-invariant (finite) result for the α^2 amplitude of Delbrück scattering in ordinary QED.

- The "infinite" renormalization of the vacuum polarization tensor then requires the following counterterm :

$$-\frac{e^2}{12\pi^2} \left[\sum_{i=0}^{i=N} c_i \log \frac{m_i^2}{m^2} \right] \left\{ \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - b^2 A_\mu A^\mu + (b_\mu A^\mu)^2 \right\} \quad (15)$$

The $\mathcal{O}(b^2)$ part of this counterterm is absent in the classical Lagrangian density, breaks gauge invariance and moreover seems to generate a b dependent mass for the photon.

- Once the previously defined "infinite" renormalization has been done, note that, contrarily to $\Pi_0^{\mu\nu}$ and $\Pi_b^{\mu\nu}$, $\Pi_{bb}^{\mu\nu}(p, m_i, b)$ does not vanish at $p^2 = 0$. Denoting as $\Pi^{(R)\mu\nu}$ the renormalized vacuum polarization tensor[§], one gets :

$$\begin{aligned} \Pi_{p^2=0}^{(R)\mu\nu} = \\ = -i \frac{e^2}{\pi^2} \int_0^1 dz \left[\frac{Z^2}{m^2} (b.p)^2 g^{\mu\nu} - \frac{Z(1-2Z)}{m^2} (b.p) (b^\mu p^\nu + b^\nu p^\mu) + \left(\frac{3Z^2}{m^2} b^2 + \frac{2Z^3}{m^4} (b.p)^2 \right) p^\mu p^\nu \right] = \end{aligned}$$

[§]Once the counterterm (15) has been added, the limit $m_i \rightarrow \infty$ may be safely taken, except for $i = 0$ where $m_0 = m$ and $c_0 = 1$

$$= -i \frac{e^2}{10m^2\pi^2} \left[\frac{1}{3} (b.p)^2 g^{\mu\nu} - (b.p)(b^\mu p^\nu + b^\nu p^\mu) + (b^2 + \frac{(p.b)^2}{7m^2}) p^\mu p^\nu \right]$$

Then, if one tried to enforce the normalization condition

$$\Pi_{p^2=0}^{(R)\mu\nu} = 0 ,$$

a necessary condition for the photon to remain massless, an extra finite counterterm would be required but, due to the last term, it would have a canonical dimension 6,

$$\frac{e^2}{70(m^2)^2\pi^2} b^\alpha b^\beta (\partial_\alpha \partial_\mu A^\mu)(\partial_\beta \partial_\nu A^\nu) \quad (16)$$

which breaks renormalizability.


All this rules out this approximation scheme and we now consider the usual "perturbative approach".

III. THE VACUUM POLARIZATION TENSOR IN EXTENDED QED: GAUGE INVARIANT APPROACH

If we no longer restrict ourselves to the linear approximation for the fermion propagator, two new Feynman graphs have to be added in the calculation of $\Pi_{bb}^{\mu\nu}(p, m, b)$:

$$\tilde{\Pi}_{bb}^{\mu\nu}(p, m, b) = e^2 \int \frac{d^4 l}{(2\pi)^4} \{ \text{Tr}[\gamma^\mu S_2(l) \gamma^\nu S_0(l+p)] + \text{Tr}[\gamma^\mu S_0(l) \gamma^\nu S_2(l+p)] \} \quad (17)$$

Diagrammatically, the new graphs are represented by (the series)



The result may be written as **

**Details on that calculation will be published elsewhere.

$$\begin{aligned}\tilde{\Pi}_{bb}^{\mu\nu}(p, m_i, b) &= \tilde{\mathbb{A}}((p.b)^2, b^2, p^2, m_i)g^{\mu\nu} + \tilde{\mathbb{B}}(p^2, m_i)b^\mu b^\nu + \\ &+ \tilde{\mathbb{C}}(p^2, m_i)(p.b)(b^\mu p^\nu + b^\nu p^\mu) + \tilde{\mathbb{D}}((p.b)^2, b^2, p^2, m_i)p^\mu p^\nu,\end{aligned}\quad (18)$$

where

$$\begin{aligned}\tilde{\mathbb{A}}((p.b)^2, b^2, p^2, m_i) &= -i\frac{e^2}{\pi^2} \left\{ -\frac{b^2}{6} \log \frac{m_i^2}{m^2} + \int_0^1 dz \left[\frac{1-8Z-8Z^2}{12\Delta_i} b^2 p^2 + \frac{4Z^2}{\Delta_i} (b.p)^2 + \right. \right. \\ &\quad \left. \left. + \frac{2Z^3}{[\Delta_i]^2} p^2 (b.p)^2 - \frac{Z^2}{2[\Delta_i]^2} (p^2)^2 b^2 \right] \right\},\end{aligned}\quad (19)$$

$$\tilde{\mathbb{B}}(p^2, m_i) = -i\frac{e^2}{\pi^2} \left\{ \frac{1}{6} \log \frac{m_i^2}{m^2} - \int_0^1 dz \left[\frac{(1-4Z)(1+2Z)}{12\Delta_i} p^2 \right] \right\}, \quad (20)$$

$$\tilde{\mathbb{C}}(p^2, m_i) = -i\frac{e^2}{\pi^2} \left\{ \int_0^1 dz \left[-\frac{2Z^2}{\Delta_i} \right] \right\}, \quad (21)$$

$$\tilde{\mathbb{D}}((p.b)^2, b^2, p^2, m_i) = -i\frac{e^2}{\pi^2} \left\{ \int_0^1 dz \left[\frac{Z(1-Z)}{2\Delta_i} b^2 + \frac{Z^2(3-4Z)}{4[\Delta_i]^2} b^2 p^2 - 2\frac{Z^3}{[\Delta_i]^2} (p.b)^2 \right] \right\}. \quad (22)$$

As in Section 2, our results were simplified thanks to integration by parts on z and the symmetry of the integral when z is changed into $(1-z)$ and we have set

$$Z = z(1-z),$$

$$\Delta_i = m_i^2 - z(1-z)p^2.$$

When these results are combined with those in Section 2, the full tensor $\bar{\Pi}_{bb}^{\mu\nu}(p, m, b)$ simplifies to :

$$\bar{\Pi}_{bb}^{\mu\nu}(p, m, b) = -i\frac{e^2}{\pi^2} \{ -X^{\mu\nu} + Y^{\mu\nu} \} \int_0^1 dz \left[\frac{Z}{\Delta_i} + \frac{Z^2}{[\Delta_i]^2} p^2 \right], \quad (23)$$

where we have introduced the two transverse tensors of dimension 4, quadratic in b^α :

$$\begin{aligned}a) X^{\mu\nu} &= b^2(g^{\mu\nu}p^2 - p^\mu p^\nu) \\ b) Y^{\mu\nu} &= g^{\mu\nu}(p.b)^2 + p^2 b^\mu b^\nu - (p.b)(p^\mu b^\nu + p^\nu b^\mu).\end{aligned}\quad (24)$$

Again, a few comments are in order:

- With that double expansion in b^α and $\hbar$, gauge invariance holds at the regularized level and then we also checked that *no ultra-violet divergence remains* in the full $\bar{\Pi}_{bb}^{\mu\nu}$.

Then the solely required "infinite" renormalization of the vacuum polarization tensor is the standard QED one

$$-\frac{e^2}{12\pi^2} \left[\sum_{i=0}^{i=N} c_i \log \frac{m_i^2}{m^2} \right] \left\{ \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right\}. \quad (25)$$

- Since $\Pi_b^{\mu\nu}$ vanishes at $p^2 = 0$, no Chern-Simons-like term appear [7].
- However, $\bar{\Pi}_{bb}^{\mu\nu}(p, m_i, b)$ still does not vanish at $p^2 = 0$. Denoting as $\bar{\Pi}^{(R)\mu\nu}$ the renormalized vacuum polarization tensor (again, once the QED counterterm has been added, the limit $m_i \rightarrow \infty$ may be taken, except for $i = 0$ where $m_0 = m$ and $c_0 = 1$), one gets:

$$\bar{\Pi}_{p^2=0}^{(R)\mu\nu} = -i \frac{e^2}{6\pi^2 m^2} [g^{\mu\nu} (p.b)^2 + b^2 p^\mu p^\nu - (p.b)(p^\mu b^\nu + p^\nu b^\mu)].$$

Then, we have two possibilities:

- in a kind of "minimal subtraction" (let us recall that the infinite subtraction has been done (25)), this non-vanishing behaviour might indicate a non vanishing mass for the photon - despite the validity of gauge invariance. In subsection 4.2, we shall analyse this possibility through the Schwinger-Dyson equation;
- in a more standard way, one may enforce the normalization condition

$$\bar{\Pi}_{p^2=0}^{(R)\mu\nu} = 0$$

in order to the photon remain massless (see subsection 4.1): then an extra finite counterterm is required. Among those possible which compensate the previous quantity up to $p^2 = 0$ vanishing contributions

$$\begin{aligned} i \frac{e^2}{6\pi^2 m^2} [g^{\mu\nu} (p.b)^2 + b^2 p^\mu p^\nu - (p.b)(p^\mu b^\nu + p^\nu b^\mu)] + \mathcal{O}(p^2) = \\ = i \frac{e^2}{6\pi^2 m^2} [Y^{\mu\nu} - X^{\mu\nu}] + \mathcal{O}'(p^2), \end{aligned}$$

we choose the unique gauge invariant term:

$$\frac{e^2}{6\pi^2 m^2} \left[g^{\mu\nu} (b^\alpha F_{\alpha\mu}) (b^\beta F_{\beta\nu}) - \frac{b^2}{4} F_{\mu\nu} F^{\mu\nu} \right]. \quad (26)$$

Let us now discuss more precisely the consequences of the Lorentz breaking on the photon mass and the Lamb-Shift.

IV. PHYSICAL CONSEQUENCES OF THE LORENTZ BREAKING

In order to obtain the full photon propagator, the Schwinger-Dyson equation must be solved [11]. In a given order of the perturbative series, it implies the summation of an infinite set of proper Feynman diagrams. In this way, as

$$(G_0^{-1})^{\mu\nu} = i[p^2 g^{\mu\nu} - (1 - \frac{1}{\alpha}) p^\mu p^\nu]$$

is the inverse of the QED free photon propagator, the Schwinger-Dyson equation ^{††}

$$(G^{-1})^{\mu\nu} = (G_0^{-1})^{\mu\nu} - \bar{\Pi}^{(R)\mu\nu}(p, m, b)$$

leads us to

$$(G^{-1})^{\mu\nu} = i \left[A' g^{\mu\nu} + B' b^\mu b^\nu + C' (b^\mu p^\nu + b^\nu p^\mu) + D' p^\mu p^\nu + E' \epsilon^{\mu\nu\alpha\beta} p_\alpha b_\beta \right], \quad (27)$$

where the values of the functions A', B', C', D' and E' depend on the finite subtraction (the normalisation condition).

From (27) we obtain the inverse tensor:

$$G_{\mu\nu} = -i \left[A'' g_{\mu\nu} + B'' b_\mu b_\nu + C'' (b_\mu p_\nu + b_\nu p_\mu) + D'' p_\mu p_\nu + E'' \epsilon_{\mu\nu\alpha\beta} p^\alpha b^\beta \right], \quad (28)$$

^{††}Recall that the renormalized tensor $\bar{\Pi}^{(R)\mu\nu}(p, m, b)$ is obtained after addition of the infinite and eventually of the finite counterterms (resp (25) and (26)): then the limit $m_i \rightarrow \infty$, $i \neq 0$, may be taken.

where^{‡‡}

$$\begin{aligned}
A'' &= \frac{A'}{Deno1}, \\
B'' &= \frac{p^2 E'^2}{A'[Deno1]} - \frac{A'B' + p^2(B'D' - C'^2)}{A'[Deno2]}, \\
C'' &= -\frac{(p.b)E'^2}{A'[Deno1]} - \frac{A'C' - (p.b)(B'D' - C'^2)}{A'[Deno2]}, \\
D'' &= \frac{b^2 E'^2}{A'[Deno1]} - \frac{A'D' + b^2(B'D' - C'^2)}{A'[Deno2]}, \\
E'' &= -\frac{E'}{Deno1},
\end{aligned} \tag{29}$$

We now discuss successively the case when a normalisation condition is imposed as well as the “minimal subtraction” scheme.

A. With the Normalization Condition $\bar{\Pi}_{p^2=0}^{(R)\mu\nu} = 0$

In this case:

$$\begin{aligned}
A'_{NC} &= p^2 \left\{ 1 + \frac{e^2}{2\pi^2} [p^2 \beta(p^2, m^2) + [(p.b)^2 - p^2 b^2] \chi(p^2, m^2)] \right\}, \\
B'_{NC} &= \frac{e^2}{2\pi^2} (p^2)^2 \chi(p^2, m^2), \\
C'_{NC} &= -\frac{e^2}{2\pi^2} (p^2) (p.b) \chi(p^2, m^2), \\
D'_{NC} &= \frac{1-\alpha}{\alpha} + \frac{e^2}{2\pi^2} [-p^2 \beta(p^2, m^2) + p^2 b^2 \chi(p^2, m^2)], \\
E'_{NC} &= -\frac{e^2}{2\pi^2} p^2 \gamma(p^2, m^2),
\end{aligned} \tag{30}$$

where:

^{‡‡}We have defined:

$$Deno1 = A'^2 + [p^2 b^2 - (p.b)^2] E'^2,$$

$$Deno2 = A'[A' + b^2 B' + 2(p.b)C' + p^2 D'] + [p^2 b^2 - (p.b)^2](B'D' - C'^2).$$

Note that if $(G^{-1})^{\mu\nu}$ were purely transverse ($A' + (p.b)C' + p^2 D' = (p.b)B' + p^2 C' = 0$), the inversion would not be possible as $Deno2 \equiv 0$.

$$\begin{aligned}
\beta(p^2, m^2) &= \int_0^1 dz \frac{[1 - 2Z - 8Z^2]}{12\Delta_0} \\
\chi(p^2, m^2) &= \frac{2}{p^2} \int_0^1 dz \left\{ \left[\frac{Z}{\Delta_0} + \frac{Z^2}{[\Delta_0]^2} p^2 \right] - \left[\frac{Z}{m^2} \right] \right\} = 2 \int_0^1 dz Z^2 \left[\frac{1}{m^2 \Delta_0} + \frac{1}{[\Delta_0]^2} \right] \\
\gamma(p^2, m^2) &= \int_0^1 dz \frac{Z}{\Delta_0}.
\end{aligned} \tag{31}$$

The functions β , χ and γ are finite at $p^2 = 0$:

$$\beta(0, m^2) = 1/(30m^2), \quad \chi(0, m^2) = 2/(15m^4), \quad \gamma(0, m^2) = 1/(6m^2).$$

We obtain:

$$\begin{aligned}
Deno1 &\simeq A'^2_{NC} \simeq (p^2)^2 \left\{ 1 + \frac{e^2}{\pi^2} [p^2 \beta(p^2, m^2) + [(p.b)^2 - p^2 b^2] \chi(p^2, m^2)] \right\} \\
Deno2 &\simeq \frac{(p^2)^2}{\alpha} \left[1 + \frac{e^2}{2\pi^2} p^2 \beta(p^2, m^2) \right].
\end{aligned}$$

As a consequence:

$$\begin{aligned}
A'' &\simeq \frac{1}{p^2} \left\{ 1 - \frac{e^2}{2\pi^2} [p^2 \beta(p^2, m^2) + [(p.b)^2 - p^2 b^2] \chi(p^2, m^2)] \right\} \\
B'' &\simeq -\frac{e^2}{2\pi^2} \chi(p^2, m^2), \\
C'' &\simeq \frac{e^2}{2\pi^2} \frac{(p.b)}{(p^2)} \chi(p^2, m^2), \\
D'' &\simeq -\frac{1}{(p^2)^2} \left\{ 1 - \alpha + \frac{e^2}{2\pi^2} [-p^2 \beta(p^2, m^2) + p^2 b^2 \chi(p^2, m^2)] \right\}, \\
E'' &\simeq \frac{e^2}{2\pi^2} \frac{1}{p^2} \gamma(p^2, m^2).
\end{aligned} \tag{32}$$

So, we have checked that, as a consequence of the normalization condition $\bar{\Pi}_{p^2=0}^{(R)\mu\nu} = 0$, the photon remains massless.

If one expands the vacuum polarisation tensor around $p^2 = 0$, one obtains a new contribution to the Lamb-shift. We consider a static configuration where $p^2 = -\vec{p}^2$ and extract from (32) an approximate expression for the Coulomb interaction:

$$\frac{e^2}{\vec{p}^2} \left[1 + \frac{e^2 \vec{p}^2}{60\pi^2 m^2} \left\{ 1 - 4 \left[\frac{(\vec{p} \cdot \vec{b})^2}{m^2 \vec{p}^2} + \frac{b^2}{m^2} \right] \right\} \right].$$

As a consequence, the effect of Lorentz breaking on that part of the Lamb-shift is drastically tiny (recall that $|b/m| < 10^{-22}$ ([14])).

B. “Minimal Subtraction” Case

If only the infinite counterterm (25) is added, one obtains:

$$\begin{aligned}
A'_{MS} &= p^2 + \frac{e^2}{2\pi^2} [(p^2)^2 \beta(p^2, m^2) + [(p.b)^2 - p^2 b^2] \phi(p^2, m^2)] , \\
B'_{MS} &= \frac{e^2}{2\pi^2} p^2 \phi(p^2, m^2), \\
C'_{MS} &= -\frac{e^2}{2\pi^2} (p.b) \phi(p^2, m^2), \\
D'_{MS} &= \frac{1-\alpha}{\alpha} + \frac{e^2}{2\pi^2} [-p^2 \beta(p^2, m^2) + b^2 \phi(p^2, m^2)], \\
E'_{MS} &= -\frac{e^2}{2\pi^2} p^2 \gamma(p^2, m^2),
\end{aligned} \tag{33}$$

where:

$$\begin{aligned}
\beta(p^2, m^2) &= \int_0^1 dz \frac{[1 - 2Z - 8Z^2]}{12\Delta_0} \\
\phi(p^2, m^2) &= 2 \int_0^1 dz \left[\frac{Z}{\Delta_0} + \frac{Z^2}{[\Delta_0]^2} p^2 \right] \\
\gamma(p^2, m^2) &= \int_0^1 dz \frac{Z}{\Delta_0}.
\end{aligned} \tag{34}$$

The functions $\beta(p^2, m^2)$, $\phi(p^2, m^2)$ and $\gamma(p^2, m^2)$ are finite at $p^2 = 0$:

$$\beta(0, m^2) = 1/(30m^2) , \quad \phi(0, m^2) = 1/(3m^2) , \quad \gamma(0, m^2) = 1/(6m^2) .$$

We now obtain:

$$\begin{aligned}
Deno1 &\simeq A'^2_{MS} \simeq p^2 \left\{ p^2 + \frac{e^2}{\pi^2} [(p^2)^2 \beta(p^2, m^2) + [(p.b)^2 - p^2 b^2] \phi(p^2, m^2)] \right\} \\
Deno2 &\simeq \frac{(p^2)^2}{\alpha} \left\{ 1 + \frac{e^2}{2\pi^2} p^2 \beta(p^2, m^2) \right\} .
\end{aligned}$$

As a consequence, a shift of the zero mass pole seems to happen. Indeed, if one expands $Deno1$ and $Deno2$ near $p^2 = 0$, one gets

$$Deno1 \simeq p^2 \left\{ p^2 + \frac{e^2}{\pi^2} [(p.b)^2 \phi(0, m^2)] + \mathcal{O}(p^2) \right\} \simeq p^2 [p^2 + \mu^2] , \quad Deno2 \simeq \frac{(p^2)^2}{\alpha}$$

with

$$\mu^2 \simeq \frac{e^2}{\pi^2} \frac{(p.b)^2}{3m^2}.$$

However, this “tachyon-like” singularity is not so simple to interpret due to its p dependence and we shall not pursue that direction.

V. CONCLUDING REMARKS

In this paper we presented two calculations for the 1-loop vacuum polarization tensor in the Lorentz and PCT violating QED defined by (1). This was done using the canonical perturbation theory and the P-V-R regularization scheme. In a first approximation we consider the electron propagator as “dressed” up to first order in b . This scheme breaks gauge invariance and we argue it should not be considered. In a second calculation, we consider the complete expansion to second order in b . Gauge invariance is then recovered and we discussed the normalisation conditions that might be added in order the photon to remain massless. Let us remark that our calculation for the $\Pi_{bb}^{\mu\nu}$ part allows us to correct a sign error in a previous calculation (formula (20) of [15]).

The full photon propagator is then obtained through the Schwinger-Dyson equation allowing a detailed analysis of its singular structure. In particular, we found that in a “minimal scheme” without normalisation condition, a new tachyonic-like pole appears. On the contrary, if the vacuum polarisation tensor is normalised such that it vanishes at $p^2 = 0$, no spurious pole appears and the photon remains massless. The effect on the Lamb-shift coming from the dependence on b of the vacuum polarisation appears to be quite negligible.

Further investigations concerning the full 1-loop renormalization of the extend QED as well as its thermal version is now in progress and will be reported in a forthcoming paper [16].

Acknowledgements. LCC and JLT are grateful to the LPTHE - Université de Paris VI et VII, where part of this work was developed. LCC acknowledges, in special, Prof. Olivier

Babelon, Director of the LPTHE, and Profs. B. Diu, B. Douçot and B. Delamotte, for the kind hospitality. LCC also thanks Fapesp-Brazil and CNRS-France for financial support. JLT thanks FAPESP-Brazil for the financial support.

REFERENCES

- [1] R. F. Streater and A. S. Wightman, “PCT, Spin, Statistics and All That”, Benjamin, NY (1964).
- [2] V. A. Kostelecký, Editor, *CPT and Lorentz Symmetry I*, World Scientific, Singapore, (1999) and, V. A. Kostelecký, Editor, *CPT and Lorentz Symmetry II*, World Scientific, Singapore (2002), and refs. therein.
- [3] S. Coleman and S. Glashow, Phys. Rev. D **59**, 116008 (1999).
- [4] R. Jackiw and V. A. Kostelecký, Phys. Rev. Lett. **82**, 3572 (1999).
- [5] A. Andrianov, P. Giacconi and R. Soldati, JHEP 0202:030 (2002).
- [6] M. Pérez-Victoria, Phys. Rev. Lett.**83**, 2518 (1999);
M. Chaichian, W. F. Chen and R. González Felipe, Phys. Lett. B **503**, 215 (2001);
J. M. Chung and B. K. Chung, Phys. Rev. D **63**, 1050015 (2001).
- [7] G. Bonneau, Nucl. Phys. B **593**, 398 (2001).
- [8] Yu. A. Sitenko and K. Yu. Rulik, Eur. Phys. J. **C28**, 405 (2003).
- [9] S. Carroll, G. Field and R. Jackiw, Phys. Rev. D **41**, 1231 (1990).
- [10] G. Rayski, Acta Phys. Polonica **9** 129 (1948);
W. Pauli and F. Villars, Rev. Mod. Phys. **21**, 434 (1949).
- [11] F. J. Dyson, Phys. Rev. **75**, 1736 (1949);
J. Schwinger, Proc. Nat. Acad. Sc., **37**, 452 e 455 (1951).
- [12] S. L. Adler, Phys. Rev. **177**, 2426 (1969);
J. S. Bell and R. Jackiw, Nuovo Cimento **60A**, 47 (1969);
T. P. Cheng and L. F. Li, *Gauge Theory of Elementary Particle Physics*, Clarendon Press, Oxford (1984).

- [13] L. C. Costa, PhD thesis, Institute of Theoretical Physics - UNESP (2004).
- [14] R. Bluhm, “QED tests of Lorentz symmetry”, arXiv:hep-ph/0411149.
- [15] W. F.Chen and G. Kunstatter, Phys.Rev. **D62**, 105029, (2000).
- [16] L. C. Costa and J. L. Tomazelli, in preparation.